

NIH-1790 Genotyping Strategies

Reaction Components	Vol (ul)
5X GoTaq Buffer	10
25mM MgCl ₂	3.5
10mM dNTPs	1
Primer 20 uM	1
Primer 20 uM	1
5 U/ul Taq polymerase	0.5
Water	28
Total mix volume	45
Tail lysate (1:20 dilution)	5
Total reaction volume	50

Step	Temp	Time	Note
1	94C	15"	
2	65C	30"	Decrease 1C/cycle
3	72C	40"	Go to 1, 10 cycles
4	94C	15"	
5	55C	30"	
6	72C	40"	Go to 4, 30 cycles

Primer Sequences (5' to 3')	
Mutant PCR: Primer LTR-2 and Primer 1790-3', 251 bp	
Recommended Wt PCR: Primer 1790-5' and Primer 1790-3', 253 bp	
Primer LTR-2	AAATGGCGTTACTTAAGCTAGCTTGC
Primer 1790-5'	CAGGATCTCACTCCGTAGCCTTGG
Primer 1790-3'	AAGGACTCCGGGAAGGTTATATGA

Well	Sample	Genotype
1	41	het
2	42	het
3	43	het
5	wt lysate	wt
6	ES DNA	het
7	water	no amp

